

Organic carbon fluxes mediated by corals at elevated $p\text{CO}_2$ and temperature

Stephen Levas^{1,6,*}, Andréa G. Grottoli¹, Mark E. Warner², Wei-Jun Cai^{2,7}, James Bauer³, Verena Schoepf^{1,8}, Justin H. Baumann¹, Yohei Matsui¹, Colin Gearing¹, Todd F. Melman⁴, Kenneth D. Hoadley², Daniel T. Pettay², Xinping Hu^{5,9}, Qian Li^{5,10}, Hui Xu^{5,11}, Yongchen Wang⁵

¹School of Earth Sciences, The Ohio State University, Columbus, OH 43210, USA

²School of Marine Science and Policy, University of Delaware, Lewes, DE 19958, USA

³Aquatic Biogeochemistry Laboratory, Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, Columbus, OH 43210, USA

⁴Reef Systems Coral Farm, New Albany, OH 43054, USA

⁵Department of Marine Sciences, University of Georgia, Athens, GA 30601, USA

⁶Present address: Department of Geography and the Environment, Villanova University, Villanova, PA 19085, USA

⁷Present address: School of Marine Science and Policy, University of Delaware, Newark, DE 19958, USA

⁸Present address: ARC Centre of Excellence for Coral Reef Studies and School of Earth and Environment, The University of Western Australia, Crawley, WA 6009, Australia

⁹Present address: Department of Physical and Environmental Sciences, Texas A&M University, Corpus Christi, TX 78412, USA

¹⁰Present address: State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 3610005, China

¹¹Present address: Department of Ocean Science and Engineering, Zhejiang University, Hangzhou 310058, China

ABSTRACT: Increasing ocean acidification (OA) and seawater temperatures pose significant threats to coral reefs globally. While the combined impacts of OA and seawater temperature on coral biology and calcification in corals have received significant study, research to date has largely neglected the individual and combined effects of OA and seawater temperature on coral-mediated organic carbon (OC) fluxes. This is of particular concern as dissolved and particulate OC (DOC and POC, respectively) represent large pools of fixed OC on coral reefs. In the present study, coral-mediated POC and DOC, and the sum of these coral-mediated flux rates (total OC, $\text{TOC} = \text{DOC} + \text{POC}$) as well as the relative contributions of each to coral metabolic demand were determined for 2 species of coral, *Acropora millepora* and *Turbinaria reniformis*, at 2 levels of $p\text{CO}_2$ (382 and 741 μatm) and seawater temperatures (26.5 and 31.0°C). Independent of temperature, DOC fluxes decreased significantly with increases in $p\text{CO}_2$ in both species, resulting in more DOC being retained by the corals and only representing between 19 and 6% of TOC fluxes for *A. millepora* and *T. reniformis*. At the same time, POC and TOC fluxes were unaffected by elevated temperature and/or $p\text{CO}_2$. These findings add to a growing body of evidence that certain species of coral may be less at risk to the impacts of OA and temperature than previously thought.

KEY WORDS: Ocean acidification · Carbon dioxide · Ocean warming · *Acropora* · *Turbinaria* · Calcification · Climate change

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Coral reefs are threatened globally due to a combination of direct and indirect human perturbations to ocean waters (e.g. Hoegh-Guldberg et al. 2007). Ris-

ing atmospheric carbon dioxide (CO_2) concentrations have contributed to increases in global seawater temperatures and an overall decrease in ocean pH (i.e. increase in ocean acidification, OA; Sabine et al. 2004, IPCC 2013). As hypercalcifying organisms that

live close to their thermal temperature limit, corals are severely threatened by both increasing temperatures and OA (Hoegh-Guldberg et al. 2007, Wild et al. 2011).

OA is the direct result of roughly half of anthropogenic CO₂ emissions being taken up by the world's oceans (Caldeira & Wickett 2003, Sabine et al. 2004). CO₂ reacts with water to form carbonic acid, which, through the production of protons (H⁺) and bicarbonate ions (HCO₃⁻), subsequently lowers pH, carbonate ion (CO₃²⁻) concentration, and the calcium carbonate saturation state (Ω) of seawater (Orr 2011). Globally, surface seawater pH has declined by an average of 0.1 units since the start of the industrial revolution (Caldeira & Wickett 2003) and is projected to decrease another 0.3 to 0.4 units by the end of this century (Orr 2011). Increasing OA is predicted to result in a decrease in coral calcification between 15 and 40% by the end of the century (Kleypas et al. 1999, 2011, Langdon et al. 2000). Under experimental OA conditions, coral calcification rates either decrease (e.g. Marubini & Atkinson 1999, Langdon & Atkinson 2005, Edmunds et al. 2012, Comeau et al. 2013b, Schoepf et al. 2013) or remain unchanged (Reynaud et al. 2003, Edmunds et al. 2012, Houlbrèque et al. 2012, Comeau et al. 2013b, Schoepf et al. 2013). In addition to calcification, OA has been shown to result in decreased tissue biomass, chlorophyll *a* (chl *a*) concentrations, and increased susceptibility to bleaching (i.e. loss of endosymbionts) in some coral species (Anthony et al. 2008, Edmunds et al. 2012, Comeau et al. 2013a), but to have limited or no effect on coral energy reserves, biomass, or chl *a* in other species (Schoepf et al. 2013).

Increased seawater temperature has also been associated with declines in coral calcification rates, tissue biomass, and energy reserves (i.e. lipids, protein, and carbohydrates) (Jokiel & Coles 1977a, Fitt et al. 1993, Edmunds et al. 2003, Marshall & Clode 2004, Grottoli et al. 2006, Rodrigues & Grottoli 2006, 2007, Cantin et al. 2010, Levas et al. 2013). When both seawater temperature and *p*CO₂ are simultaneously elevated, calcification either decreases (Reynaud et al. 2003, Anthony et al. 2008, Rodolfo-Metalpa et al. 2011) or remains unchanged (Langdon & Atkinson 2005, Rodolfo-Metalpa et al. 2010, Edmunds 2011, Schoepf et al. 2013). In the case of the 2 species used in the present study (*Acropora millepora* and *Turbinaria reniformis*), calcification rates do not change, energy reserves do not decline, and photosynthesis and/or algal density decline or do not change when both seawater temperature and *p*CO₂ are simultaneously elevated (Schoepf et al. 2013).

However, despite the implications for coral physiology and coral reef biogeochemical carbon (C) cycling, no studies to date have assessed the individual or interactive effects of elevated temperatures and *p*CO₂ on coral-mediated organic matter fluxes. This is of particular concern, as corals play an important role in coral reef biogeochemical cycles and fauna–microbe interactions through the production of mucus (Wild et al. 2004a,b, 2005a,b, 2008, 2009) and its subsequent degradation into particulate and dissolved organic matter (Brown & Bythell 2005, Bythell & Wild 2011).

Coral-mediated particulate and dissolved organic matter may have unique ecological functions in coral reef ecosystems. In healthy coral reefs, corals secrete particulate organic matter in the form of mucus that acts as a trap for organic and inorganic matter which they can ingest or slough off, where it may be deposited to the seafloor, where roughly 90% of coral-mediated particulate organic C (POC) is degraded by benthic microbial consumers (Wild et al. 2004a). Meanwhile, coral-mediated production of dissolved organic C (DOC) is rapidly utilized and supports heterotrophic microbes in the water column (Nelson et al. 2013). In addition to producing POC and DOC, corals have been shown to take up DOC and POC under normal and elevated temperature conditions, presumably as a source of fixed C (Ferrier-Pagès et al. 1998, Anthony 2000, Haas et al. 2010, Naumann et al. 2010, Tremblay et al. 2012, Levas et al. 2013), which may serve as a supplemental nutritional resource under stressful conditions and be associated with coral resilience. Under elevated temperatures, coral bleaching results in decreases in the acquisition of photosynthetically fixed C and often results in the depletion of energy reserves (Fitt et al. 1993, Edmunds et al. 2003, Grottoli et al. 2006, Rodrigues & Grottoli 2007). At the same time, OA potentially makes calcification more energetically costly (Cohen & Holcomb 2009, Erez et al. 2011, Pandolfi et al. 2011). Therefore, under stressful environmental conditions such as elevated temperatures and OA, coral fixed C budgets could be dramatically strained. Thus, we hypothesize that some corals may alter their DOC and POC fluxes in order to conserve organic carbon (OC) as a mechanism to tolerate the stress and/or promote recovery. This has implications for coral reef biogeochemical cycles and host–microbe interactions through the production of mucus, which supports a diverse microbial community (Wild et al. 2004a,b, 2005a,b, 2008, 2009), its subsequent degradation into POC and DOC (Brown &

Bythell 2005, Bythell & Wild 2011), and ensuing fueling of the reef microbial loop.

In the present study, we evaluated the individual and interactive effects of seawater $p\text{CO}_2$ and temperature on coral-mediated POC and DOC fluxes, total OC (TOC = DOC + POC) fluxes, and the relative contributions of each flux to coral metabolic demand in 2 species of Pacific coral. Collectively, these measurements were used to evaluate how coral-mediated OC fluxes may change with the predicted increases in baseline seawater temperature and seawater $p\text{CO}_2$ forecasted for the end of this century.

MATERIALS AND METHODS

Experimental design

Coral collection, acclimation, experimental design, and procedures used in the present study and the seawater chemical parameters (Table 1) have been described previously by Schoepf et al. (2013). Briefly, 6 parent colonies each of *Acropora millepora* and *Turbinaria reniformis* were collected at 3 to 10 m depth in northwest Fiji (17° 29' 19" S, 177° 23' 39" E) in April 2011 and transported to Reef Systems Coral Farm in New Albany, Ohio, USA. Four coral fragments were collected from each parent colony (total of 24 fragments per species), mounted onto pre-labeled tiles, and allowed to acclimate in a holding tank with natural light for 2.5 mo.

In July 2011, all 48 fragments were transferred to experimental recirculating indoor aquaria with custom-made artificial seawater (ESV Aquarium Products), which was designed to mimic the chemical composition and alkalinity of natural reef seawater. Fragments were then allowed to acclimate for another 10 d at ambient seawater temperature and $p\text{CO}_2$ (26.5°C and $p\text{CO}_2$ of 382 μatm) and light (275 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ on a 9:15 h light:dark cycle) prior to exposure to the experimental conditions.

Four experimental treatments were used: 26.5°C with 382 μatm $p\text{CO}_2$, 26.5°C with 741 μatm $p\text{CO}_2$, 31.0°C with 382 μatm $p\text{CO}_2$, and 31.0°C with 741 μatm $p\text{CO}_2$. The 2 $p\text{CO}_2$ levels used (382 and 741 μatm) were designed to represent present-day $p\text{CO}_2$ and the average $p\text{CO}_2$ level expected by the second half of the 21st century (IPCC 2013). The control temperature (26.5°C) represents the current average annual temperatures in Fiji (www.ospo.noaa.gov/Products/ocean/index.html). Temperatures in the elevated seawater temperature treatments were increased gradually over the course of the study in order to avoid shocking the corals (Fig. 1). Each seawater temperature– $p\text{CO}_2$ combination consisted of a separate recirculating tank system with one 905 l sump and 6 aquaria of 57 l each. One fragment from each parent colony per species was placed in 1 of the 6 aquaria in each system such that each parent colony of each species was represented in each system. The experiment lasted for 24 d. Throughout the experiment, corals were fed every 3 d with 48 h old brine shrimp nauplii. Each fragment was also buoy-




Fig. 1. Mean \pm SE daily temperature for each treatment throughout the study. Errors are sometimes very small, making the error bars smaller than the symbol and not visible

Table 1. Mean \pm SE seawater conditions for each treatment representing 2 $p\text{CO}_2$ levels and 2 temperature regimes as reported by Schoepf et al. (2013). Temp.: temperature; T: total scale; TA: total alkalinity; Ω_{arag} : saturation state of aragonite

Variable	Ambient $p\text{CO}_2$		Elevated $p\text{CO}_2$	
	Ambient temp.	Elevated temp.	Ambient temp.	Elevated temp.
Temperature (°C)	26.45 \pm 0.01	29.31 \pm 0.02	26.61 \pm 0.01	28.93 \pm 0.02
pH_T	8.07 \pm 0.01	8.04 \pm 0.01	7.83 \pm 0.01	7.81 \pm 0.01
$p\text{CO}_2$ (μatm)	364.31 \pm 9.69	400.62 \pm 16.83	732.04 \pm 22.37	749.63 \pm 26.21
TA ($\mu\text{mol kg}^{-1}$)	2269.9 \pm 10.84	2270.1 \pm 11.15	2306.3 \pm 10.64	2304.5 \pm 9.081
Ω_{arag}	3.69 \pm 0.07	3.79 \pm 0.09	2.40 \pm 0.06	2.52 \pm 0.06

antly weighed on Days 1, 12, and 24 of the experiment, and calcification rates per cm² were determined for the first and the second half of the experiment as reported by Schoepf et al. (2013).

POC, DOC, and TOC fluxes

To assess coral-mediated organic matter fluxes, individual coral fragments were incubated in circular 1.25 l UV-transparent acrylic chambers (15.3 cm tall × 10.7 cm diameter). Since these incubations were only conducted during the day, calculated values reflect daytime fluxes. Six coral fragments of the same species and treatment were cleaned of any epizootic organisms and detritus and their tiles thoroughly cleaned of encrusting organisms and algae. Each fragment was transferred to an individual chamber freshly filled with treatment-specific seawater, with air exposure limited to ~1 s. A lid was securely placed on top of the chamber, making sure to eliminate all air bubbles, and the chamber was fully submerged in a large tank supplied with treatment-specific flowing seawater to maintain seawater temperature. A seventh chamber contained no coral fragments and served as a blank control.

Water circulation inside each chamber was maintained using a Teflon stir-bar. Dissolved oxygen and seawater temperature inside each chamber were monitored with an YSI 550A submersible polarographic probe that was inserted into the base of each chamber and O-ring sealed. Duplicate 1 l seawater samples, representing the initial seawater conditions, were collected from the treatment-specific flowing seawater supply line into 2 acid-cleaned 1 l polycarbonate brown bottles and placed on ice.

After 1.5 h, the water level in the incubation tank was lowered to expose the top 3 cm of the chambers. All coral fragments were then removed and placed back in the respective treatment aquaria, and the seawater from each chamber was collected into an individual acid-cleaned 1 l brown bottle and placed on ice. The incubations were limited to 1.5 h to ensure that dissolved oxygen did not reach supersaturation, which can be a potential stressor. Each 1 l sample was then filtered onto a pre-combusted (500°C for 4 h) GF/F filter (0.7 µm nominal pore size) and stored at –20°C for POC analysis. A 30 ml subsample of the filtrate was collected in an acid-cleaned polycarbonate bottle, acidified to a pH ≤2 using ultra high purity HCl, and stored at –20°C for DOC analysis. Duplicate GF/F filters and 30 ml aliquots were also collected from the blank chamber

for POC and DOC analysis, respectively. Two sets of incubations on separate fragments were performed per day (one starting at 10:00 h and the other starting at 14:00 h), and all incubations were completed over the course of the last 4 d of the experiment. To facilitate comparisons among treatments within a species, *A. millepora* incubations were always conducted from 10:00 to 11:30 h and *T. reniformis* incubations were always conducted from 14:00 to 15:30 h. Chambers were cleaned and thoroughly rinsed with treatment-appropriate seawater prior to each incubation.

DOC concentrations were determined using high-temperature catalytic oxidation using a Shimadzu model 5050 TOC analyzer. After thawing, samples were sparged 2 min with ultra-high purity air, and then 250 µl of sample were injected into the instrument. A standard consisting of 1000 µM glucose was used to establish a 4-point calibration curve. A regular quality control check consisting of glucose was also injected after every 5 samples. The standard deviation of replicate measurements of the glucose standard was ±4%.

For POC analyses, GF/F filters were dried for 24 h at 60°C and then acidified in a clean glass desiccator using concentrated fuming HCl for 24 h to remove inorganic C. POC was determined using a Costech Elemental Analyzer (EA). An acetanilide standard was used to establish a 4-point calibration curve and as a quality control standard spaced throughout each EA sample run. The standard deviation of the replicate measurements of the standard was ±1.8%.

For each coral incubation, the DOC and POC concentrations were corrected for potential microheterotrophic and microautotrophic biases by subtracting the POC and DOC concentrations of their blank chamber from their respective values. Since microbially mediated DOC degradation is not observed in the first 2 h (Nakajima et al. 2009), a short incubation time combined with the use of a blank correction minimizes any interference of bacterial degradation in the signal. Fluxes (i.e. release or uptake; µmol C h⁻¹ cm⁻²) were calculated as the difference between the blank-corrected OC concentration (µmol) of the coral fragment and the initial seawater concentration, and standardized to the incubation duration (1.5 h) and fragment surface area (FSA; cm²) as determined by the foil technique (Marsh 1970) using the following equation:

$$\text{Daytime OC flux} = \frac{(\text{OC}_{\text{coral}} - \text{OC}_{\text{blank}}) - \text{OC}_{\text{Initial}}}{1.5 \times \text{FSA}} \quad (1)$$

Negative fluxes indicate a net uptake by coral fragments, and positive fluxes indicate a net release from

coral fragments. TOC flux was calculated as the sum of the individual POC and DOC fluxes.

Proportionate contribution of POC, DOC, and TOC to coral respiration

The proportionate contribution of heterotrophically derived OC to the coral daytime C budget was calculated relative to respiration, which is termed the percent contribution of heterotrophy to animal respiration (CHAR; Grottoli et al. 2006). Here, CHAR for POC (CHAR_{POC}), DOC (CHAR_{DOC}), and TOC (CHAR_{TOC}) were determined as follows.

Respiration was measured on each coral fragment individually during the last 4 d of the experiment using previously described methods (Fitt & Warner 1995). For a given fragment, respiration incubations were conducted 1 d before the POC and DOC incubations in order to minimize any potential stress to the corals and to ensure that both sets of measurements were contemporaneous. In brief, coral fragments were individually placed in 350 ml water-tight UV-transparent acrylic chambers filled with their respective treatment seawater and a stir bar, and dissolved gaseous oxygen was lowered to 80 % by bubbling with ultra high purity nitrogen gas. Chambers were then placed into a 20 l Plexiglas tank filled with fresh water, which was temperature controlled by a heater/chiller unit and set to 26.5°C or 31.0°C, depending on the treatment. The incubation tank was then covered by black plastic, and dissolved oxygen was measured for approximately 15 min. Respiration was then determined by calculating the slope of the oxygen concentration recorded during the last 10 min of each run.

Daytime CHAR_{POC} and CHAR_{DOC} for each fragment were calculated as the daytime flux of POC or DOC in $\mu\text{g C}$ standardized to grams ash-free dry weight h^{-1} (POC_f and DOC_f , respectively), divided by the $\mu\text{g C}$ lost via daytime respiration h^{-1} (R_c), assuming a mole-to-mole relationship of O_2 consumed to CO_2 produced during respiration. Thus, daytime CHAR_{POC} and CHAR_{DOC} were calculated as:

$$\text{CHAR}_{\text{POC}} = \frac{\text{POC}_f}{R_c} \times 100\% \quad (2)$$

$$\text{CHAR}_{\text{DOC}} = \frac{\text{DOC}_f}{R_c} \times 100\% \quad (3)$$

Therefore here, CHAR_{POC} and CHAR_{DOC} are the percentage of a coral's respiration that can be met or lost through OC uptake or released during the day.

CHAR_{TOC} for each fragment was calculated as the sum of CHAR_{POC} and CHAR_{DOC} . Negative CHAR values indicate a net loss of OC relative to respiratory demand, and positive CHAR values indicate a net gain of OC relative to respiratory demand. We recognize that POC and DOC fluxes (S. Levas unpubl. data) and respiration (Rodrigues & Grottoli 2007) vary between day and night. Since we only have daytime measurements for all 3 variables, our interpretation is limited to daytime fluxes.

Statistical analysis

Three-way mixed-model ANOVA was run for each species separately and tested the effects of temperature, $p\text{CO}_2$, and genotype on POC fluxes, DOC fluxes, TOC fluxes, daytime respiration, CHAR_{POC} , CHAR_{DOC} , and CHAR_{TOC} . Temperature and $p\text{CO}_2$ were fixed and fully crossed, whereas genotype was a random factor. The purpose of including the genotype in the ANOVA model was to determine whether any single genotype was systematically different from all others for a given variable. In cases where significant genotype effects were detected, Tukey tests revealed that all average values overlapped among genotypes. As such, we concluded that the selected colonies represented the natural variation in the population, and no single or group of colonies was consistently different from the others. Thus, interaction terms involving genotype were not included in the ANOVAs. All data were deemed to be normally distributed using Shapiro-Wilk normality tests. All data sets met the assumptions of homogeneity of variances as determined with plots of expected versus residual value. Tukey post hoc tests were performed when main effects were significant ($p \leq 0.05$). *A posteriori* slice tests (e.g. tests of simple effects, Winer 1971) determined whether low (382 μatm) and high (741 μatm) $p\text{CO}_2$ treatment averages significantly differed within each temperature treatment. Bonferroni corrections were not applied due to increased likelihood of false negatives (Quinn & Keough 2002, Moran 2003). To explore possible relationships between a suite of published variables on these same corals and the data in this study, Pearson correlation analyses were run between calcification rates and total energy reserve data from Schoepf et al. (2013) and POC, DOC, and TOC fluxes here. Statistical analyses were performed using SAS software, Version 9.3 of the SAS System for Windows.

RESULTS

POC, DOC, TOC fluxes, and daytime respiration

In *Acropora millepora*, POC flux rates were unaffected by both temperature and $p\text{CO}_2$ (Fig. 2A, Table 2). However, DOC fluxes averaged 59% lower in the elevated $p\text{CO}_2$ treatments (741 μatm) than in ambient $p\text{CO}_2$ treatments (382 μatm ; Fig. 2B, Table 2). This $p\text{CO}_2$ effect was most pronounced at ambient temperature (Fig. 2B). No significant temperature or $p\text{CO}_2$ effect was detected for *A. millepora* TOC fluxes (Fig. 2C, Table 2), while daytime respiration rates were 39% lower in elevated $p\text{CO}_2$ treatments than in ambient conditions irrespective of temperature (Fig. 2D, Table 2). Overall, DOC fluxes constituted about 45% of the TOC flux at ambient $p\text{CO}_2$ and 19% at elevated $p\text{CO}_2$ (Fig. 2). *Turbinaria*




Fig. 2. Mean \pm SE daytime particulate, dissolved, and total organic carbon (POC, DOC, and TOC, respectively) fluxes and daytime respiration for (A–D) *Acropora millepora* and (E–H) *Turbinaria reniformis*. Within each panel, asterisks indicate significant differences between 382 and 741 μatm averages within a specific temperature (determined by a *posteriori* slice tests). Positive rates indicate release/loss, whereas negative rates indicate uptake. Sample sizes ranged between 5 and 6 coral fragments in each treatment

reniformis POC fluxes were unaffected by both temperature and $p\text{CO}_2$, while DOC fluxes decreased by an average of 100% under elevated $p\text{CO}_2$, independent of temperature (Fig. 2E,F, Table 2). Despite significant declines in DOC fluxes at elevated $p\text{CO}_2$,

Table 2. Results of 3-way ANOVAs for *Acropora millepora* and *Turbinaria reniformis* particulate organic carbon (POC), dissolved organic carbon (DOC), and total organic carbon (TOC) fluxes, and daytime respiration at 2 temperatures (26.5°C and 31.0°C) and 2 $p\text{CO}_2$ levels (382 and 741 μatm). Effects of temperature (T) and $p\text{CO}_2$ were fixed and fully crossed. Genotype (G) was a random effect. Significant effects are shown in **bold**

Variable	Effect	df	F	p
A. millepora				
POC	Model	8,22	0.78	0.6308
	T	1	0.28	0.6035
	$p\text{CO}_2$	1	0.85	0.3708
	G	5	1.04	0.4339
	$T \times p\text{CO}_2$	1	0.01	0.9375
DOC	Model	8,21	3.10	0.0343
	T	1	1.33	0.2961
	$p\text{CO}_2$	1	10.62	0.0062
	G	5	1.64	0.2194
	$T \times p\text{CO}_2$	1	2.60	0.1311
TOC	Model	8,22	1.11	0.4145
	T	1	0.97	0.3426
	$p\text{CO}_2$	1	0.00	0.9958
	G	5	1.57	0.2308
	$T \times p\text{CO}_2$	1	0.08	0.7805
Daytime respiration	Model	8,22	1.78	0.1655
	T	1	0.01	0.9338
	$p\text{CO}_2$	1	5.86	0.0297
	G	5	1.03	0.4381
	$T \times p\text{CO}_2$	1	2.69	0.1231
T. reniformis				
POC	Model	8,23	0.72	0.6709
	T	1	0.04	0.8397
	$p\text{CO}_2$	1	0.38	0.5474
	G	5	0.90	0.5034
	$T \times p\text{CO}_2$	1	0.83	0.3774
DOC	Model	8,21	3.68	0.0185
	T	1	0.72	0.4119
	$p\text{CO}_2$	1	21.21	0.0005
	G	5	1.17	0.3749
	$T \times p\text{CO}_2$	1	2.16	0.1656
TOC	Model	8,21	0.61	0.7586
	T	1	0.00	0.9904
	$p\text{CO}_2$	1	0.02	0.8997
	G	5	0.68	0.6445
	$T \times p\text{CO}_2$	1	1.30	0.2747
Daytime respiration	Model	8,21	1.42	0.2643
	T	1	10.09	0.0063
	$p\text{CO}_2$	1	0.04	0.8506
	G	5	0.25	0.9346
	$T \times p\text{CO}_2$	1	0.03	0.8730

there were no overall temperature or $p\text{CO}_2$ effects on *T. reniformis* TOC fluxes (Fig. 2G, Table 2). However, *T. reniformis* daytime respiration rates were 40% greater at elevated temperatures but unaffected by elevated $p\text{CO}_2$ (Fig. 2H, Table 2). Overall, DOC fluxes constituted about 30% of the TOC flux at ambient $p\text{CO}_2$ and 6% at elevated $p\text{CO}_2$ (Fig. 2).

Proportionate contribution of POC, DOC, and TOC to coral respiration

A. millepora CHAR_{POC} , CHAR_{DOC} , and CHAR_{TOC} as well as *T. reniformis* CHAR_{POC} and CHAR_{TOC} were unaffected by temperature and $p\text{CO}_2$ (Fig. 3A–D, F, Table 3). However, in *T. reniformis*, CHAR_{DOC} was more negative at ambient compared to elevated $p\text{CO}_2$ levels, and positive at elevated temperature




Fig. 3. Mean \pm SE daytime percent contribution of heterotrophically derived particulate, dissolved, and total organic carbon (POC, DOC, and TOC, respectively) to animal respiration (CHAR_{POC} , CHAR_{DOC} , and CHAR_{TOC} , respectively) for (A–C) *Acropora millepora* and (D–F) *Turbinaria reniformis* in each treatment. Within each panel, asterisks indicate significant differences between 382 and 741 μatm averages within a specific temperature (determined by a *posteriori* slice tests). Negative values indicate carbon losses. Sample sizes ranged between 5 and 6 individual coral fragments in each treatment

(Fig. 3E, Table 3). Overall, CHAR_{DOC} constituted 31% and 25% of the CHAR_{TOC} at ambient and elevated $p\text{CO}_2$ for *A. millepora*, respectively, and 26% and –1% of the CHAR_{TOC} at ambient and elevated $p\text{CO}_2$ for *T. reniformis*, respectively (Fig. 3).

Relationship of POC, DOC, and TOC to calcification and energy reserves

A. millepora POC and TOC fluxes were not significantly correlated with respiration, calcification rates, or total energy reserves (Table 4). Only *A. millepora*

Table 3. Results of 3-way ANOVAs for *Acropora millepora* and *Turbinaria reniformis*, daytime percent contribution of heterotrophically derived particulate, dissolved, and total organic carbon (POC, DOC, and TOC, respectively) to animal respiration (CHAR_{POC} , CHAR_{DOC} , and CHAR_{TOC} , respectively) at 2 temperatures (26.5°C and 31.0°C) and 2 $p\text{CO}_2$ levels (382 and 741 μatm). Effects of temperature (T) and $p\text{CO}_2$ were fixed and fully crossed. Genotype (G) was a random effect. Significant effects are shown in **bold**

Variable	Effect	df	F	p
<i>A. millepora</i>				
CHAR_{POC}	Model	8,2	1.62	0.2062
	T	1	0.09	0.7732
	$p\text{CO}_2$	1	3.34	0.0889
	G	5	2.00	0.1403
	$T \times p\text{CO}_2$	1	0.01	0.9412
CHAR_{DOC}	Model	8,22	1.65	0.2040
	T	1	1.87	0.1943
	$p\text{CO}_2$	1	2.87	0.1140
	G	5	1.27	0.3332
	$T \times p\text{CO}_2$	1	0.90	0.3609
CHAR_{TOC}	Model	8,21	1.76	0.1760
	T	1	0.89	0.3639
	$p\text{CO}_2$	1	1.40	0.2579
	G	5	2.43	0.0916
	$T \times p\text{CO}_2$	1	0.00	0.9609
<i>T. reniformis</i>				
CHAR_{POC}	Model	8,21	0.92	0.5286
	T	1	1.67	0.2174
	$p\text{CO}_2$	1	1.47	0.2457
	G	5	0.63	0.6792
	$T \times p\text{CO}_2$	1	1.02	0.3303
CHAR_{DOC}	Model	8,21	1.96	0.0298
	T	1	1.27	0.2810
	$p\text{CO}_2$	1	17.0	0.0012
	G	5	1.28	0.3305
	$T \times p\text{CO}_2$	1	3.61	0.0800
CHAR_{TOC}	Model	8,20	0.71	0.6768
	T	1	1.24	0.2857
	$p\text{CO}_2$	1	0.45	0.5155
	G	5	0.51	0.7645
	$T \times p\text{CO}_2$	1	1.45	0.2495

Table 4. Pearson's correlation analyses for *Acropora millepora* and *Turbinaria reniformis*. POC: particulate organic carbon flux ($\mu\text{mol C h}^{-1} \text{cm}^{-2}$); DOC: dissolved organic carbon flux ($\mu\text{mol C h}^{-1} \text{cm}^{-2}$); TOC: total organic carbon flux (DOC + POC; $\mu\text{mol C h}^{-1} \text{cm}^{-2}$); Day R: daytime respiration ($\mu\text{mol C h}^{-1} \text{cm}^{-2}$); Calc.1: calcification during the first 12 d of the experiment ($\text{mg d}^{-1} \text{cm}^{-2}$); Calc.2: calcification during the last 12 d of the experiment ($\text{mg d}^{-1} \text{cm}^{-2}$); E. reserves: sum of protein + carbohydrate + lipid concentrations (mg g^{-1} ash-free dry weight). Calcification and E. reserve data from Schoepf et al. (2013). Significant correlations are shown in **bold**

Variable		<i>A. millepora</i>				<i>T. reniformis</i>			
		Day R	Calc.1	Calc.2	E. reserves	Day R	Calc.1	Calc.2	E. reserves
POC	r	0.096	0.143	-0.039	-0.150	0.152	0.194	0.201	0.201
	p	0.65	0.51	0.86	0.49	0.47	0.36	0.34	0.34
	N	23	23	22	23	24	24	24	24
DOC	r	-0.369	0.055	0.537	-0.395	-0.269	-0.007	-0.072	-0.350
	p	0.09	0.81	0.01	0.06	0.22	0.97	0.74	0.10
	N	22	21	21	22	22	22	22	22
TOC	r	-0.017	0.185	0.166	-0.275	0.158	0.272	0.238	0.145
	p	0.93	0.40	0.45	0.20	0.48	0.22	0.28	0.51
	N	23	23	22	23	22	22	22	22

DOC fluxes were positively correlated with calcification rates during the second half of the experiment (Table 4). *T. reniformis* POC, DOC, and TOC fluxes were not significantly correlated with respiration, calcification rates, or total energy reserves (Table 4).

DISCUSSION

Previous studies have investigated OC fluxes by corals and coral reef organisms under ambient (Crossland et al. 1980, Crossland 1987, Tanaka et al. 2008, 2009, Haas et al. 2010, Naumann et al. 2010) and varying environmental conditions such as eutrophication (Tanaka et al. 2007), elevated temperature (Haas et al. 2010, Levas et al. 2013), and during different seasons (Naumann et al. 2010). However, this study is the first to assess the combined impacts of elevated $p\text{CO}_2$ and temperature on coral-mediated OC fluxes.

POC, DOC, and TOC fluxes

Both coral species had positive POC fluxes (i.e. released POC) irrespective of $p\text{CO}_2$ or temperature treatments (Fig. 2A,E). Average POC fluxes for *Acropora millepora* of 3.36 to 6.08 $\text{mg POC m}^{-2} \text{h}^{-1}$ were within the range of those reported for *Acropora* corals from Japan and the Red Sea of 1.0 to 7.1 $\text{mg POC m}^{-2} \text{h}^{-1}$ (Tanaka et al. 2008, 2009, Naumann et al. 2010). *Turbinaria reniformis* POC fluxes in this

study varied from 8.80 to 18.24 $\text{mg POC m}^{-2} \text{h}^{-1}$ and are similar to other coral species from the Great Barrier Reef (Wild et al. 2005b). The lack of any elevated temperature effect on either species of coral contrasts with previous studies showing that corals have higher POC fluxes during the summer months compared to the winter months (Naumann et al. 2010) and when bleached (Niggel et al. 2008). In addition, we show that neither elevated temperature nor $p\text{CO}_2$ altered coral-mediated POC fluxes or the corresponding CHAR_{POC} (Figs. 2A,E & 3A,D). As most coral-mediated POC is likely derived from mucus, this would suggest that elevated $p\text{CO}_2$ —unlike other environmental stressors such as increased sedimentation, light availability, and temperature (reviewed by Bythell & Wild 2011)—has no effect on coral mucus production. Thus, coral POC fluxes appear to be unaffected by the treatment conditions imposed in this study.

While OA had no effect on *A. millepora* and *T. reniformis* POC fluxes, both species had reduced DOC fluxes under increased acidification irrespective of temperature (Fig. 2B,F). Here, DOC fluxes for *A. millepora* and *T. reniformis* ranged from 1.17 to 3.72 $\text{mg DOC m}^{-2} \text{h}^{-1}$ and -0.53 to 5.09 $\text{mg DOC m}^{-2} \text{h}^{-1}$, respectively. This is within the range of DOC flux measurements reported for healthy *Acropora* ranging from 1.9 $\text{mg DOC m}^{-2} \text{h}^{-1}$ (Tanaka et al. 2008) to 30.7 $\text{mg DOC m}^{-2} \text{h}^{-1}$ (Naumann et al. 2010). The lack of any significant temperature effect on DOC fluxes is consistent with previous findings for *A. millepora* and *T. reniformis* (Niggel et al. 2008). However, previous measurements on temperature stres-

sed *Porites* corals showed an increase in DOC flux rates (Haas et al. 2010), which suggests that there may be location- or population-specific responses with the same species. We hypothesize as to the mechanisms driving the retention or release of DOC under elevated $p\text{CO}_2$ below, but further study is needed to fully test these ideas.

Interestingly, despite the decrease in DOC fluxes for all *A. millepora* under elevated $p\text{CO}_2$, there was no corresponding decrease in CHAR_{DOC} because respiration decreased under all conditions (Figs. 2D & 3B). At the same time, *A. millepora* maintained energy reserves while calcification declined (Schoepf et al. 2013). Together, these findings suggest that this coral species either modifies its DOC losses or is incapable of maintaining its DOC fluxes. Either way, limiting the loss of organic C via DOC fluxes does not prevent calcification from declining under $p\text{CO}_2$ stress conditions (Schoepf et al. 2013). Decreases in DOC fluxes at elevated $p\text{CO}_2$ were even more pronounced in *T. reniformis* (Fig. 2E), with dramatic decreases in CHAR_{DOC} when both $p\text{CO}_2$ and temperature were elevated (Fig. 3E). At the same time, *T. reniformis* maintained energy reserves and calcification rates (Schoepf et al. 2013). Together these findings show that this coral species is also capable of modifying its DOC losses, and suggests that DOC modification may play a role in optimizing the coral's C-budget and maintaining calcification under OA and elevated temperature conditions.

Net TOC fluxes (POC flux + DOC flux) were always positive (i.e. corals released TOC) and did not significantly vary with temperature and/or $p\text{CO}_2$ (Fig. 2C, G). TOC fluxes ranged from 5.5 to 7.8 $\text{mg C m}^{-2} \text{h}^{-1}$ for *A. millepora* and from 10.8 to 20.0 $\text{mg C m}^{-2} \text{h}^{-1}$ for *T. reniformis* (Fig. 2C,G). These values are up to 9 times higher than net TOC fluxes by control and temperature-stressed *Stylophora pistillata* (Tremblay et al. 2012). Thus, TOC fluxes appear to vary widely among species. In addition, CHAR_{TOC} was not significantly affected by elevated temperature and/or $p\text{CO}_2$ (Fig. 3C,F), despite the significant decline in CHAR_{DOC} in *T. reniformis*. This is because TOC is predominantly composed of POC, which was not affected by the treatment conditions. Furthermore, C lost as TOC accounted for 14 to 47% of coral metabolic demand (Fig. 3C,F). This is in contrast to *S. pistillata*, which has positive CHAR_{TOC} when healthy and only small OC losses (i.e. CHAR_{TOC} losses of 3 to 6%) when temperature stressed (Tremblay et al. 2012). This would suggest that OC uptake and/or production is highly species and health-status specific.

Model for coral OC loss under normal and elevated $p\text{CO}_2$

Corals release POC in the form of mucus that coats the outer epidermal surface of the animal and as DOC exudates (Fig. 4, adapted from Brown & Bythell




Fig. 4. Model of coral mucus as particulate organic carbon (POC) and dissolved organic carbon (DOC) release in (A) healthy and (B) ocean acidification (OA) stressed corals. In healthy corals, carbon translocation associated with POC and DOC production is as follows (solid arrows in A). (1) Endosymbiotic algae (algae) photosynthetically fix carbon and translocate it to the coral host. (2) Fixed carbon can also be acquired from heterotrophic consumption of plankton. (3) Fixed carbon is either secreted as DOC to the water column or is translocated to mucus glands. (4) Mucus glands secrete DOC and (5) mucus. (6) The sloughed mucus constitutes the coral holobiont-mediated POC, which can also (7) dissolve or be bacterially degraded to form DOC (adapted from Brown & Bythell 2005). In OA stressed corals, pathways 3, 4, and 7 are diminished (dashed lines)

2005). In healthy corals, endosymbionts translocate excess photosynthetically fixed C to the host (Muscatine 1967, 1990, Patton et al. 1977, Harland et al. 1991). While some of this fixed C is stored as energy reserves (Battey & Patton 1984, Edmunds & Spencer Davies 1986), any excess is either released directly to the water column as DOC or transferred to the mucocytes (mucus glands) and excreted as mucus (Crossland et al. 1980) (Fig. 4A). Much of this mucus is sloughed off of the coral surface and while some remains as POC, up to 80% ultimately ends up in the DOC pool (Wild et al. 2004b, Nakajima et al. 2009). In addition, heterotrophically acquired C also contributes fixed C that can contribute to coral-produced mucus and DOC (Ferrier-Pagès et al. 1998).

However, under OA conditions, decreased DOC fluxes, as observed for both *A. millepora* and *T. reniformis*, could impact this model of C-transport in 3 possible ways (Fig. 4B): (1) decreases in the secretion of fixed C in the form of DOC directly to the water column (arrow 3 in Fig. 4), (2) decreases in the secretion of DOC from the mucocytes (arrow 4), and/or (3) decreases in the breakdown of the mucus POC to DOC (arrow 7). Neither coral decreased its POC production, which would suggest that the breakdown of POC into DOC would remain unchanged, thus the third pathway is not likely involved in the reduction of DOC fluxes. Rather, both coral species appear to be actively conserving C by decreasing direct losses of DOC from the first and/or second mechanism (Fig. 4B) in response to OA.

Corals may not limit POC loss as mucus due to the important physio-chemical role that mucus plays in protecting corals from environmental conditions such as increased sedimentation, temperature, and UV exposure (reviewed by Brown & Bythell 2005). Limiting mucus production and subsequent sloughing of mucus could increase susceptibility to environmental perturbations and potentially pathogens as well (Ducklow & Mitchell 1979, Rublee et al. 1980). However, a large fraction of DOC is comprised of lipid (Crossland 1987) which represents a potential energy source. Thus, while POC fluxes are not altered, limiting the loss of DOC through mechanisms (1) and (2) therefore has the potential to limit the amount of lipids lost in the form of DOC. This hypothesis is consistent with recent findings showing that lipid concentrations did not decrease in either species when exposed to elevated $p\text{CO}_2$ (Schoepf et al. 2013). It also indicates that corals can be flexible in their organic matter fluxes and respiration rates in such a way that maintains overall TOC irrespective of OA and/or temperature stress.

Implications

Our findings indicate that coral organic matter fluxes are highly variable and largely species-specific in their responses to elevated $p\text{CO}_2$ and temperature. Decreases in DOC fluxes were accompanied by decreases in respiration resulting in stable CHAR_{TOC} . In addition, retaining DOC has the potential to help sustain energy reserves. This is consistent with previous findings showing that biomass and energy reserves are maintained in both *A. millepora* and *T. reniformis* (Schoepf et al. 2013). It is also consistent with emerging findings that increased $p\text{CO}_2$ does not cause significant additional energetic demand on calcification (Edmunds 2011, 2012, McCulloch et al. 2012, Schoepf et al. 2013). However, since DOC represents a significant source of reduced C for planktonic bacteria (Ducklow & Mitchell 1979), decreases in coral-mediated DOC released to the water column could negatively impact reef microbial communities. Overall, these findings add to a growing body of evidence that certain species of corals may be less at risk to the impacts of OA and temperature than previously thought (Edmunds 2012, McCulloch et al. 2012, Schoepf et al. 2013).

It is imperative to keep in mind that these results are based on a short-term (24 d) exposure to elevated temperature and $p\text{CO}_2$. As described by Comeau et al. (2014), it is ideal for experiments testing the impacts of OA to last months to years in order to allow the organisms to acclimate. Clearly, further studies are needed to assess how coral DOC and POC fluxes respond to elevated temperature and $p\text{CO}_2$ in other species, and over longer periods of time. Longer studies are essential to determine whether changes in coral DOC and POC production ultimately affect coral C budgets and calcification in the long term. In addition, further research is needed to explore how microbial community composition and density might change under future global change scenarios and how that might influence coral OC fluxes.

Acknowledgements. We thank E. Zebrowski, M. Berzelis, M. Ringwald, A. Barrett, and the staff at Reef Systems Coral Farm for their field and laboratory support. This work was supported by funding from the National Science Foundation (grants NSF-EF-1041124, 1040940, 1041070). We thank 3 anonymous reviewers for their comments and suggestions on a previous version of this paper. Data have been deposited at the Biological & Chemical Oceanography Data Management Office and are available at www.bco-dmo.org/project/528004.

LITERATURE CITED

- Anthony KRN (2000) Enhanced particle-feeding capacity of corals. *Coral Reefs* 19:59–67
- Anthony KRN, Kline DI, Diaz-Pulido G, Dove S, Hoegh-Guldberg O (2008) Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proc Natl Acad Sci USA* 105:17442–17446
- Batley JF, Patton JS (1984) A reevaluation of the role of glycerol in carbon translocation in zooxanthellae–coelenterate symbiosis. *Mar Biol* 79:27–38
- Brown BE, Bythell JC (2005) Perspectives on mucus secretion in reef corals. *Mar Ecol Prog Ser* 296:291–309
- Bythell JC, Wild C (2011) Biology and ecology of coral mucus release. *J Exp Mar Biol Ecol* 408:88–93
- Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. *Nature* 425:365
- Cantin NE, Cohen AL, Karnauskas KB, Tarrant AM, McCorkle DC (2010) Ocean warming slows coral growth in the central Red Sea. *Science* 329:322–325
- Cohen AL, Holcomb M (2009) Why corals care about ocean acidification—uncovering the mechanism. *Oceanography* 22:118–127
- Comeau S, Carpenter R, Edmunds P (2013a) Effects of feeding and light intensity on the response of the coral *Porites rus* to ocean acidification. *Mar Biol* 160:1127–1134
- Comeau S, Edmunds PJ, Spindel NB, Carpenter RC (2013b) The responses of eight coral reef calcifiers to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnol Oceanogr* 58:388–398
- Comeau S, Edmunds PJ, Spindel NB, Carpenter RC (2014) Fast coral reef calcifiers are more sensitive to ocean acidification in short-term laboratory incubations. *Limnol Oceanogr* 59:1081–1091
- Crossland CJ (1987) In situ release of mucus and DOC-lipid from the corals *Acropora variabilis* and *Stylophora pistillata* in different light regimes. *Coral Reefs* 6:35–42
- Crossland CJ, Barnes DJ, Borowitzka MA (1980) Diurnal lipid and mucus production in the staghorn coral *Acropora acuminata*. *Mar Biol* 60:81–90
- Ducklow HW, Mitchell R (1979) Composition of mucus released by coral-reef coelenterates. *Limnol Oceanogr* 24:706–714
- Edmunds PJ (2011) Zooplanktivory ameliorates the effects of ocean acidification on the reef coral *Porites* spp. *Limnol Oceanogr* 56:2402–2410
- Edmunds PJ (2012) Effect of pCO₂ on the growth, respiration, and photophysiology of massive *Porites* spp. in Moorea, French Polynesia. *Mar Biol* 159:2149–2160
- Edmunds PJ, Spencer Davies P (1986) An energy budget for *Porites porites* (Scleractinia). *Mar Biol* 92:339–347
- Edmunds PJ, Gates RD, Gleason DF (2003) The tissue composition of *Montastraea franksi* during a natural bleaching event in the Florida Keys. *Coral Reefs* 22:54–62
- Edmunds PJ, Brown D, Moriarty V (2012) Interactive effects of ocean acidification and temperature on two scleractinian corals from Moorea, French Polynesia. *Glob Change Biol* 18:2173–2183
- Erez J, Reynaud S, Silverman J, Schneider K, Allemand D (2011) Coral calcification under ocean acidification and global change. In: Stambler N, Dubinsky Z (eds) *Coral reefs: an ecosystem in transition*. Springer, New York, NY, p 151–176
- Ferrier-Pagès C, Gattuso JP, Cauwet G, Jaubert J, Allemand D (1998) Release of dissolved organic carbon and nitrogen by the zooxanthellate coral *Galaxea fascicularis*. *Mar Ecol Prog Ser* 172:265–274
- Fitt WK, Warner ME (1995) Bleaching patterns of four species of Caribbean reef corals. *Biol Bull (Woods Hole)* 189:298–307
- Fitt WK, Spero HJ, Halas J, White MW, Porter JW (1993) Recovery of the coral *Montastrea annularis* in the Florida Keys after the 1987 Caribbean 'bleaching event'. *Coral Reefs* 12:57–64
- Grottoli AG, Rodrigues LJ, Palardy JE (2006) Heterotrophic plasticity and resilience in bleached corals. *Nature* 440:1186–1189
- Haas AF, Jantzen C, Naumann MS, Iglesias-Prieto R, Wild C (2010) Organic matter release by the dominant primary producers in a Caribbean reef lagoon: implication for *in situ* O₂ availability. *Mar Ecol Prog Ser* 409:27–39
- Harland AD, Fixter LM, Spencer Davies P, Anderson RA (1991) Distribution of lipids between the zooxanthellae and animal compartment in the symbiotic sea anemone *Anemonia viridis*: wax esters, triglycerides and fatty acids. *Mar Biol* 110:13–19
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS and others (2007) Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737–1742
- Houlbrèque F, Rodolfo-Metalpa R, Jeffree R, Oberhänsli F and others (2012) Effects of increased pCO₂ on zinc uptake and calcification in the tropical coral *Stylophora pistillata*. *Coral Reefs* 31:101–109
- IPCC (Intergovernmental Panel on Climate Change) (2013) *Climate change 2013: the physical science basis. Summary for policymakers. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Available at www.ipcc.ch
- Jokiel PL, Coles SL (1977a) Effects of temperature on the mortality and growth of Hawaiian reef corals. *Mar Biol* 43:201–208
- Kleypas JA, Buddemeier RW, Archer D, Gattuso JP, Langdon C, Opdyke BN (1999) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284:118–120
- Kleypas JA, Anthony KRN, Gattuso JP (2011) Coral reefs modify their seawater carbon chemistry—case study from a barrier reef (Moorea, French Polynesia). *Glob Change Biol* 17:3667–3678
- Langdon C, Atkinson MJ (2005) Effect of elevated pCO₂ on photosynthesis and calcification on corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *J Geophys Res* 110:C09S07, doi:10.1029/2004JC002576
- Langdon C, Takahashi T, Sweeney C, Chipman D and others (2000) Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochem Cycles* 14:639–654, doi:10.1029/1999GB001195
- Levas SJ, Grottoli AG, Hughes A, Osburn CL, Matsui Y (2013) Physiological and biogeochemical traits of bleaching and recovery in the mounding species of coral *Porites lobata*: implications for resilience in mounding corals. *PLoS ONE* 8:e63267
- Marsh JA (1970) Primary productivity of reef-building calcareous red algae. *Ecology* 51:255–263
- Marshall AT, Clode P (2004) Calcification rate and the effect of temperature in a zooxanthellate and an azooxanthellate scleractinian reef coral. *Coral Reefs* 23:218–224

- Marubini F, Atkinson MJ (1999) Effects of lowered pH and elevated nitrate on coral calcification. *Mar Ecol Prog Ser* 188:117–121
- McCulloch M, Falter J, Trotter J, Montagna P (2012) Coral resilience to ocean acidification and global warming through pH up-regulation. *Nat Clim Change* 2:623–633
- Moran MD (2003) Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos* 100:403–405
- Muscatine L (1967) Glycerol excretion by symbiotic algae from corals and *Tridacna* and its control by the host. *Science* 156:516–519
- Muscatine L (1990) The role of symbiotic algae in carbon and energy flux in reef corals. In: Dubinsky Z (ed) *Ecosystems of the world. Book 25: Coral reefs*. Elsevier, New York, NY, p 75–87
- Nakajima R, Yoshida T, Azman BAR, Zaleha K, Othman BHR, Toda T (2009) *In situ* release of coral mucus by *Acropora* and its influence on the heterotrophic bacteria. *Aquat Ecol* 43:815–823
- Naumann MS, Haas A, Struck U, Mayr C, El-Zibdah M, Wild C (2010) Organic matter release by dominant hermatypic corals of the Northern Red Sea. *Coral Reefs* 29:649–659
- Nelson CE, Goldberg SJ, Kelly LW, Haas AF, Smith JE, Rohwer F, Carlson CA (2013) Coral and macroalgal exudates vary in neutral sugar composition and differentially enrich reef bacterioplankton lineages. *ISME J* 7: 962–979
- Niggel W, Glas M, Laforsch C, Mayr C, Wild C (2008) First evidence of coral bleaching stimulating organic matter release by reef corals. *Proc 11th Int Coral Reef Symp, Ft. Lauderdale, FL* 2:905–911
- Orr JC (2011) Recent and future changes in ocean carbonate chemistry. In: Gattuso JP, Hansson L (eds) *Ocean acidification*. Oxford University Press, Oxford, p 41–66
- Pandolfi JM, Connolly SR, Marshall DJ, Cohen AL (2011) Projecting coral reef futures under global warming and ocean acidification. *Science* 333:418–422
- Patton JS, Abraham S, Benson AA (1977) Lipogenesis in the intact coral *Pocillopora capitata* and its isolated zooxanthellae: evidence for a light-driven carbon cycle between symbiont and host. *Mar Biol* 44:235–247
- Quinn GP, Keough MJ (2002) *Experimental design and data analysis for biologists*. Cambridge University Press, New York, NY
- Reynaud S, Leclercq N, Romaine-Lioud S, Ferrier-Pagès C, Jaubert J, Gattuso JP (2003) Interacting effects of CO₂ partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Glob Change Biol* 9: 1660–1668
- Rodolfo-Metalpa R, Martin S, Ferrier-Pagès C, Gattuso JP (2010) Response of the temperate coral *Cladocora caespitosa* to mid- and long-term exposure to pCO₂ and temperature levels projected for the year 2100 AD. *Biogeosciences* 7:289–300
- Rodolfo-Metalpa R, Houlbrèque F, Tambutté E, Boisson F and others (2011) Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat Clim Change* 1:308–312
- Rodrigues LJ, Grottoli AG (2006) Calcification rate and the stable carbon, oxygen, and nitrogen isotopes in the skeleton, host tissue, and zooxanthellae of bleached and recovering Hawaiian corals. *Geochim Cosmochim Acta* 70:2781–2789
- Rodrigues LJ, Grottoli AG (2007) Energy reserves and metabolism as indicators of coral recovery from bleaching. *Limnol Oceanogr* 52:1874–1882
- Rublee PA, Lasker HR, Gottfried M, Roman MR (1980) Production and bacterial colonization of mucus from the soft coral *Briarum asbestinum*. *Bull Mar Sci* 30:888–893
- Sabine CL, Feely RF, Gruber N, Key RM and others (2004) The oceanic sink for anthropogenic CO₂. *Science* 305: 367–371
- Schoepf V, Grottoli AG, Warner ME, Cai WJ and others (2013) Coral energy reserves and calcification in a high-CO₂ world at two temperatures. *PLoS ONE* 8:e.0075049
- Tanaka Y, Miyajima T, Koike I, Hayashibara T, Ogawa H (2007) Imbalanced coral growth between organic tissue and carbonate skeleton caused by nutrient enrichment. *Limnol Oceanogr* 52:1139–1146
- Tanaka Y, Miyajima T, Koike I, Hayashibara T, Ogawa H (2008) Production of dissolved and particulate organic matter by the reef-building corals *Porites cylindrica* and *Acropora pulchra*. *Bull Mar Sci* 82:237–245
- Tanaka T, Miyajima T, Umezawa Y, Hyashibara T, Ogawa H, Koike I (2009) Net release of dissolved organic matter by the scleractinian coral *Acropora pulchra*. *J Exp Mar Biol Ecol* 377:101–106
- Tremblay P, Naumann MS, Sikorski S, Grover R, Ferrier-Pagès C (2012) Experimental assessment of organic carbon fluxes in the scleractinian coral *Stylophora pistillata* during a thermal and photo stress event. *Mar Ecol Prog Ser* 453:63–77
- Wild C, Huettel M, Klueter A, Kremb SG, Rasheed MYM, Jørgensen BB (2004a) Coral mucus functions as an energy carrier and particle trap in the reef ecosystem. *Nature* 428:66–70
- Wild C, Rasheed M, Werner U, Franke U, Johnstone R, Huettel M (2004b) Degradation and mineralization of coral mucus in reef environments. *Mar Ecol Prog Ser* 267: 159–171
- Wild C, Rasheed M, Jantzen C, Cook P, Struck U, Huettel M, Boetius A (2005a) Benthic metabolism and degradation of natural particulate organic matter in carbonate and silicate reef sands of the northern Red Sea. *Mar Ecol Prog Ser* 298:69–78
- Wild C, Woyt H, Huettel M (2005b) Influence of coral mucus on nutrient fluxes in carbonate sands. *Mar Ecol Prog Ser* 287:87–98
- Wild C, Mayr C, Wehrmann L, Schöttner S, Naumann M, Hoffmann F, Rapp HT (2008) Organic matter release by cold water corals and its implication for fauna-microbe interaction. *Mar Ecol Prog Ser* 372:67–75
- Wild C, Wehrmann LM, Mayr C, Schöttner SI, Allers E, Lundälv T (2009) Microbial degradation of cold-water coral-derived organic matter: potential implication for organic C cycling in the water column above Tisler Reef. *Aquat Biol* 7:71–80
- Wild C, Hoegh-Guldberg O, Naumann MS, Colombo-Pallotta MF and others (2011) Climate change impedes scleractinian corals as primary reef ecosystem engineers. *Mar Freshw Res* 62:205–215
- Winer BJ (1971) *Statistical principles in experimental design*. McGraw-Hill, New York, NY